

Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

Forename(s)

Candidate signature

A-level MATHEMATICS

Unit Further Pure 2

Friday 24 June 2016

Morning

Time allowed: 1 hour 30 minutes

Materials

For this paper you must have:

- the blue AQA booklet of formulae and statistical tables.

You may use a graphics calculator.

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- Write the question part reference (eg (a), (b)(i) etc) in the left-hand margin.
- You must answer each question in the space provided for that question. If you require extra space, use an AQA supplementary answer book; do **not** use the space provided for a different question.
- Do not write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 75.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.



Answer **all** questions.

Answer each question in the space provided for that question.

1 (a) Given that $f(r) = \frac{1}{4r-1}$, show that

$$f(r) - f(r+1) = \frac{A}{(4r-1)(4r+3)}$$

where A is an integer.

[2 marks]

(b) Use the method of differences to find the value of $\sum_{r=1}^{50} \frac{1}{(4r-1)(4r+3)}$, giving your answer as a fraction in its simplest form.

[4 marks]

QUESTION
PART
REFERENCE

Answer space for question 1



3 The arc of the curve with equation $y = 4 - \ln(1 - x^2)$ from $x = 0$ to $x = \frac{3}{4}$ has length s .

(a) Show that $s = \int_0^{\frac{3}{4}} \left(\frac{1 + x^2}{1 - x^2} \right) dx$.

[4 marks]

(b) Find the value of s , giving your answer in the form $p + \ln N$, where p is a rational number and N is an integer.

[6 marks]

QUESTION
PART
REFERENCE

Answer space for question 3



4 (a) Given that $y = \tan^{-1} \sqrt{3x}$, find $\frac{dy}{dx}$, giving your answer in terms of x .

[2 marks]

(b) Hence, or otherwise, show that $\int_{\frac{1}{3}}^1 \frac{1}{(1+3x)\sqrt{x}} dx = \frac{\sqrt{3}\pi}{n}$, where n is an integer.


[4 marks]

QUESTION
PART
REFERENCE

Answer space for question 4



- 5 (a)** Find the modulus of the complex number $-4\sqrt{3} + 4i$, giving your answer as an integer. **[2 marks]**
- (b)** The locus of points, L , satisfies the equation $|z + 4\sqrt{3} - 4i| = 4$.
- (i)** Sketch the locus L on the Argand diagram below. **[3 marks]**
- (ii)** The complex number w lies on L so that $-\pi < \arg w \leq \pi$.
Find the least possible value of $\arg w$, giving your answer in terms of π . **[2 marks]**
- (c)** Solve the equation $z^3 = -4\sqrt{3} + 4i$, giving your answers in the form $re^{i\theta}$, where $r > 0$ and $-\pi < \theta \leq \pi$. **[5 marks]**

QUESTION
PART
REFERENCE**Answer space for question 5****(b)(i)**

6 (a) Given that $y = \sinh x$, use the definition of $\sinh x$ in terms of e^x and e^{-x} to show that $x = \ln(y + \sqrt{y^2 + 1})$.

[4 marks]

(b) A curve has equation $y = 6 \cosh^2 x + 5 \sinh x$.

(i) Show that the curve has a single stationary point and find its x -coordinate, giving your answer in the form $\ln p$, where p is a rational number.

[5 marks]

(ii) The curve lies entirely above the x -axis. The region bounded by the curve, the coordinate axes and the line $x = \cosh^{-1} 2$ has area A .

Show that

$$A = a \cosh^{-1} 2 + b\sqrt{3} + c$$

where a , b and c are integers.

[5 marks]

QUESTION
PART
REFERENCE

Answer space for question 6



8 (a) By applying de Moivre's theorem to $(\cos \theta + i \sin \theta)^4$, where $\cos \theta \neq 0$, show that

$$(1 + i \tan \theta)^4 + (1 - i \tan \theta)^4 = \frac{2 \cos 4\theta}{\cos^4 \theta}$$

[3 marks]

(b) Hence show that $z = i \tan \frac{\pi}{8}$ satisfies the equation $(1 + z)^4 + (1 - z)^4 = 0$, and express the three other roots of this equation in the form $i \tan \phi$, where $0 < \phi < \pi$.

[2 marks]

(c) Use the results from part **(b)** to find the values of:

(i) $\tan^2 \frac{\pi}{8} \tan^2 \frac{3\pi}{8}$;

[4 marks]

(ii) $\tan^2 \frac{\pi}{8} + \tan^2 \frac{3\pi}{8}$.

[4 marks]

QUESTION
PART
REFERENCE

Answer space for question 8



There are no questions printed on this page

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Copyright information

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2016 AQA and its licensors. All rights reserved.

